

AD-A148 811


DIOXODODECAISOPROPOXYTETRATUNGSTEN OXYGEN ATOM  
ABSTRACTION FROM ACETONE I. (U) INDIANA UNIV AT  
BLOOMINGTON DEPT OF CHEMISTRY T P BLATCHFORD ET AL.  
29 NOV 84 INDU/DC/TR-84/2-MC F/G 7/2

1/1

UNCLASSIFIED

NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

12

AD-A148 811

OFFICE OF NAVAL RESEARCH

Contract No. N00014-79-C-0044

Task No. NR 056-703

TECHNICAL REPORT NO. INDU/DC/TR-84/2-MC

DIOXODODECAISOPROPOXYTETRATUNGSTEN. OXYGEN ATOM ABSTRACTION  
FROM ACETONE IN REACTIONS WITH HEXAISOPROPOXYDITUNGSTEN ( $M \equiv M$ )

by

Timothy P. Blatchford, Malcolm H. Chisholm, Kirsten Folting

and John C. Huffman

prepared for publication

in

Chemical Communications

Department of Chemistry  
Indiana University  
Bloomington, IN 47405

November 29, 1984

Reproduction in whole or in part is permitted for  
any purpose of the United States Government.

This document has been approved for public release  
and sale; its distribution is unlimited.

DTIC  
ELECTE  
S DEC 28 1984 D  
E

DTIC FILE COPY

84 12 17 006

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER INDU/DC/TR-83/2-MC	2. GOVT ACCESSION NO. AD-A148811	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Dioxododecaisopropoxytetratungsten. Oxygen Atom Abstraction from Acetone in Reactions with Hexaisopropoxyditungsten (M-M)		5. TYPE OF REPORT & PERIOD COVERED Technical Report 1984
7. AUTHOR(s) Timothy F. Blatchford, Malcolm H. Chisholm, Kirsten Folting and John C. Huffman		6. PERFORMING ORG. REPORT NUMBER INDU/DC/TR-84/2-MC
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Indiana University Bloomington, IN 47405		8. CONTRACT OR GRANT NUMBER(s) N00C14-79-C-0044
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE November 29, 1984
		13. NUMBER OF PAGES 7
		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Hexaisopropoxyditungsten (M-M Triple bond)		
18. SUPPLEMENTARY NOTES		Accession For NTIS GRA&I <input checked="" type="checkbox"/> DTIC TAB <input type="checkbox"/> Unannounced <input type="checkbox"/> Justification By _____ Distribution/ Availability Codes
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) tungsten, alkoxides, oxo, clusters, McMurray Reagent		Dist Avail and/or Special A-1
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) W <sub>2</sub> (OPr <sup>i</sup> ) <sub>6</sub> (py) <sub>2</sub> (M) and acetone (1 equiv) react in hydrocarbon solvents and ambient temperatures to yield tetramethylethylene and W <sub>4</sub> O <sub>7</sub> (OPr <sup>i</sup> ) <sub>12</sub> which is shown, by x-ray study, to be an unusual cluster type in which the 8 electrons available for M-M bonding are partitioned to a M-M triple and M-M single bond. Dioxododecaisopropoxytetratungsten		

Dioxododecaisopropoxytungsten. Oxygen Atom Abstraction  
from Acetone in Reactions with Hexaisopropoxyditungsten (M≡M)

by Timothy P. Blatchford, Malcolm H. Chisholm\*, Kirsten  
Folting and John C. Huffman

(Department of Chemistry and Molecular Structure Center,  
Indiana University, Bloomington, Indiana 47405, USA)

Summary  $W_2(OPr^i)_6(py)_2(M \equiv M)$  and acetone (1 equiv) react in hydrocarbon solvents and ambient temperatures to yield tetramethylethylene and  $W_4O_2(OPr^i)_{12}$  which is shown, by an x-ray study, to be an unusual cluster type in which the 8 electrons available for M-M bonding are partitioned to a M-M triple and M-M single bond.

Since their discovery the hexaalkoxides of dimolybdenum<sup>1,2</sup> and ditungsten<sup>3,4</sup> have provided entry into a rich field of reaction chemistry.<sup>5</sup> The metal-metal triple bonds provide a source of electrons for redox reactions and the dinuclear center may provide a template for the assembly of substrate molecules or serve to obviate the need for consecutive reduction steps. The reduction of  $C \equiv C$ ,  $C \equiv N$  and  $C=O$  bonds in the reactions involving  $W_2(OR)_6$  compounds exemplify these principles.<sup>5</sup> We describe here a reaction between  $W_2(OPr^i)_6(py)_2$  and acetone which, by oxygen atom abstraction, leads to a novel tetranuclear cluster  $W_4O_2(OPr^i)_{12}$ .

Hydrocarbon solutions (hexane, benzene or toluene) of  $W_2(OPr^i)_6(py)_2$  and acetone (1 equiv) react<sup>†</sup> according to eq. 1 at ambient temperatures.



Analyses of the volatiles of the reaction by  $^1\text{H}$  NMR spectroscopy and g.c./m.s. showed only pyridine, tetramethylethylene and traces of acetone and isopropanol. The tungsten containing residue was obtained as black crystals from toluene and was shown to be  $\text{W}_4\text{O}_2(\text{OPr}^i)_{12}$  by single crystal x-ray studies. The central  $\text{W}_4\text{O}_{14}$  skeleton of the molecule is shown in Figure 1.

Though molybdenum and tungsten are known to form a wide variety of tetranuclear clusters including  $\text{M}_4$  tetrahedra,<sup>6</sup> butterflies,<sup>7,8</sup> squares,<sup>7</sup> rectangles,<sup>9</sup> parallelograms<sup>10,11</sup> and even chains,<sup>12</sup> the  $\text{W}_4$  skeleton seen here is unique. The average oxidation state of tungsten is +4 yielding 8 electrons for M-M bonding. Based on W-W distances,<sup>5,13</sup> these are evidently used to form one W-W triple bond,  $\text{W}(1)-\text{W}(2) = 2.404(2) \text{ \AA}$ , and one W-W single bond,  $\text{W}(3)-\text{W}(4) = 2.684(2) \text{ \AA}$ . Any direct M-M bonding at the distance of 2.95  $\text{\AA}$  must be weak by comparison. The molecule may thus be viewed as a dimer of dinuclear species. Note the  $\text{W}=\text{W}$  bond is unbridged by any atom and the two tungsten atoms,  $\text{W}(1)$  and  $\text{W}(2)$ , are 3 and 4 coordinate, respectively. The latter observation is very unusual and has only been seen recently in  $(\text{Pr}^i\text{O})_3\text{Mo}=\text{Mo}(\text{OPr}^i)(\text{CH}_2\text{Ph})_2(\text{PMe}_3)$ .<sup>14</sup> The oxo bridges may be partitioned:  $\text{O}(5)$  as  $\text{O}^-$  to  $\text{W}(1)$ , i.e.  $\text{O}(5)$  is like an alkoxy group in which  $\text{W}(3)$  substitutes for a carbon atom, and  $\text{O}(6)$  as an oxygen dative bond to  $\text{W}(2)$ . Consistent with this line of reasoning  $\text{W}(3)-\text{O}(5) = 1.86(2)$  and  $\text{W}(3)-\text{O}(6) = 2.01(1) \text{ \AA}$  representing formally W-O double and single (dative) bond distances, respectively. The other half of the molecule involving  $\text{W}(3)$  and  $\text{W}(4)$  is a confacial bioctahedron,  $(\text{RO})(\text{O}^-)(\text{O}^{2-})\text{W}(\mu\text{-OR})_3\text{W}(\text{OR})_3$ , containing a  $\text{W}_2^{10+}$  core:(M-M).

The oxygen atom abstraction reaction and coupling of ketonic carbon atoms in eq. 1 has an obvious parallel with the McMurray reagent<sup>15</sup> which employs reduced titanium, probably finely divided Ti metal or at least clusters.<sup>16</sup> In the present instance it is probable that C-C bond formation occurs to give a diolate, akin to the reactions reported by Cotton and Walton and their co-workers<sup>17</sup> involving W=W bonds. From here the reaction could proceed to give C-O bond cleavage and the alkene followed by cluster formation or alternatively the reaction with  $W_2(OPr^i)_6$ , as a reducing agent, could give a  $W_4$ -cluster from which alkene is released.

Further studies aimed at elucidating the mechanism of this reaction and evaluating its scope toward organic syntheses are planned.<sup>18</sup>

---

<sup>†</sup>Dry and oxygen free atmospheres ( $N_2$ ) and solvents were used throughout.

<sup>†</sup>Crystal data for  $W_4O_2(OPr^i)_{12}$  at  $-140^\circ C$ :  $a = 13.386(7)$  Å,  $b = 19.426(15)$  Å,  $c = 10.250(6)$  Å,  $\alpha = 99.28(4)^\circ$ ,  $\beta = 104.20(3)^\circ$ ,  $\gamma = 94.52(3)^\circ$ ,  $Z = 2$ ,  $d_{calcd} = 1.938$  g cm<sup>-3</sup> and space group  $\bar{P}1$ . Of the 6658 unique reflections collected using Mo  $K_\alpha$ ,  $6^\circ \leq 2\theta \leq 45^\circ$ , the 5423 having  $F > 3\sigma(F)$  were used in the full matrix least-squares refinement, using anisotropic thermal parameters on the W atoms, while O and C atoms were isotropic. The hydrogen atoms were located in fixed positions. No absorption or extinction corrections were used. There was no evidence of solvent molecules being present. Final residuals are  $R(F) = 0.065$  and  $R_w(F) = 0.065$ . The atomic coordinates for this work are available on request from the Director of the Cambridge Crystallographic Data Centre, University Chemical

Laboratory, Lensfield Road, Cambridge CB2 1EW. Any request should be accompanied by the full literature citation for this communication. The complete structure report, MSC No. 84036, is available from the Indiana University Library in Microfiche form only.

1. M.H. Chisholm and W.W. Reichert, J. Am. Chem. Soc., 1974, 96, 1249.
2. M.H. Chisholm, F.A. Cotton, C.A. Murillo and W.W. Reichert, Inorg. Chem., 1977, 16, 1801.
3. M.H. Chisholm and M. Extine, J. Am. Chem. Soc., 1975, 97, 5625.
4. M. Akiyama, M.H. Chisholm, F.A. Cotton, M.W. Extine, D.A. Haitko, D. Little and P.E. Fanwick, Inorg. Chem., 1979, 18, 2266.
5. M.H. Chisholm, Polyhedron, 1983, 2, 681 and references cited therein.
6. C. Perrin, R.Chevrel and M. Sergent, C.R. Hebd. Seances Acad. Sci. Ser. C., 1975, 280, 949.
7. M.H. Chisholm, R.J. Errington, K. Folting and J.C. Huffman, J. Am. Chem. Soc., 1982, 104, 2025.
8. S. Stersrad, B.J. Helland, M.W. Babich, R.A. Jacobson and R.E. McCarley, J. Am. Chem. Soc., 1978, 100, 6257.
9. R.N. McGinnis, T.R. Ryan and R.E. McCarley, J. Am. Chem. Soc., 1978, 100, 7900.
10. M.H. Chisholm, J.C. Huffman, C.C. Kirkpatrick, J. Leonelli and K. Folting, J. Am. Chem. Soc., 1981, 103, 6093.
11. R.E. McCarley, M.H. Luly, R.T. Ryan and C.C. Torardi, ACS Symp. Ser., 1981, 155, 41.




12. M. Akiyama, M.H. Chisholm, F.A. Cotton, M.W. Extine, D.A. Haitko, J. Leonelli and D. Little, J. Am. Chem. Soc., 1981, 101, 779.
13. F.A. Cotton and R.A. Walton in 'Multiple Bonds Between Metal Atoms', Wiley, 1982.
14. M.H. Chisholm, J.C. Huffman and R.J. Tatz, J. Am. Chem. Soc., submitted.
15. J.E. McMurray, Acc. Chem. Res., 1983, 16, 405.
16. R. Dams, M. Malinowski, I. Westdorp and H.Y. Geise, J. Org. Chem., 1982, 47, 248.
17. F.A. Cotton, D. DeMarco, L.R. Falvello and R.A. Walton, J. Am. Chem. Soc., 1982, 104, 7375.
18. We thank the Office of Naval Research and the Wrubel Computing Center for support of this work.

Caption to Figure

Figure 1.

The central  $W_4O_{14}$  skeleton of the  $W_4O_{12}(OPr^i)_{12}$  molecule. O(5) and O(6) are oxo oxygen atoms. Some pertinent distances (Å) and angles ( $^\circ$ ) are: W(1)-W(2) = 2.404(2), W(1)-W(3) = 2.948(2); W(2)-W(3) = 2.950(2); W(3)-W(4) = 2.684(2); W(1)-O(6), -O(7), -O(8) = 1.845(13), 1.865(14), 1.856(13); W(2)-O(5), -O(9), -O(10), -O(11) = 1.942(13), 1.910(16), 1.883(17), 1.951(16); W(3)-O(5), -O(6), -O(12), -O(13), -O(14), -O(15) = 1.857(13), 2.008(13), 1.902(14), 2.197(13), 2.125(14), 2.055(13); W(4)-O(13), -O(14), -O(15), -O(16), -O(17), -O(18) = 2.016(13), 2.054(14), 2.054(13), 1.888(17), 1.910(15), 1.805(14); W(2)-O(5)-W(3) = 101.9(6), W(1)-O(6)-W(3) = 99.8(6); W(2)-W(1)-O(6) = 108.1(4); W(2)-W(1)-O(7) = 105.2(4), W(2)-W(1)-O(8) = 106.9(4); W(1)-W(2)-O(5) = 103.9(4); W(1)-W(2)-O(9) = 101.1(5); W(1)-W(2)-O(10) = 99.9(5); W(1)-W(2)-O(11) = 102.5(5); W(1)-W(2)-W(3) = 65.91(5); W(2)-W(1)-W(3) = 65.98(5); W(1)-W(3)-W(2) = 48.11(4); W(1)-W(3)-W(4) = 137.48(5); W(2)-W(3)-W(4) = 131.50(5).



**END**

**FILMED**

1-85

**DTIC**